

EVALUATION OF BIOLOGICAL ACTIVE FORMULATES AND STRAINS FOR THE BIOLOGICAL CONTROL OF REPLANT DISEASE IN CITRUS ROOTSTOCKS

Hasan Pinar ^{1,*}, Suat Kaymak ¹, Aydin Uzun ¹

¹ Erciyes University, Department of Horticulture, Kayseri, Turkey



Abstract

Knowledge of biotic components of soil “replant disorders” or “replant disease” affecting fruit tree orchards substantially increased in the 1980s, thanks to the in-studies carried out on soil fumigants. In fact, the significant increase of young fruit tree plant growth in fumigated plots in comparison to non-fumigated plots suggested that soil microorganisms play an important role in growth reduction generally classified as “replant disorders”. The replant problems affecting all fruit tree crops of high economic value in Europe are strongly related to land degradation problems due to intensive cropping with frequent orchard replants. The combination of the selected resources with low input agro-management practices for controlling soil borne pathogens, such as solarisation, bio-pesticides, soil movement and other available tools etc., should lead to new management options in organic fruit tree orchards. In present study was conducted to determine effect of some product of Bio-formulates in citrus plants which protect from replant disease such as *Phytophthora citrophthora*. Sour orange and Trifliage rootstocks were used and treated ten active ingredient/microorganism for *Phytophthora citrophthora*. All products had positive effect on plant growing, but ISR200 had highest value. This differences can be because of plant material, temperature and product doses.

Keywords: bio-formulates, Citrus, rootstocks, replant diseases

1. INTRODUCTION

The EU 27 trend of fruit production has decreased from 70 to 60 million tons during the period from 2000 to 2009 (European Fruit Summit, 2010). There are many causes of this yield decline, however one of the main reasons is that of replant disorders due to the eco-functional intensification of growing areas specialising in fruit production occurring in Europe (Savory, 1966; Manici et al., 2003), as wells as in other parts of the world (Mazzola, 1998; van Schoor et al., 2009; Gordon Braun et al., 2010).

Knowledge of biotic components of soil “replant disorders” or “replant disease” affecting fruit tree orchards substantially increased in the 1980s, thanks to the in-studies carried out on soil fumigants. In fact, the significant increase of young fruit tree plant growth in fumigated plots in comparison to non-fumigated plots suggested that soil microorganisms play an important role in growth reduction generally classified as “replant disorders” (Browne et al., 2006). The main agents of replant disease are several soilborne fungal pathogens which can vary according to soil type, cropping practices, crops etc. They are generally defined as “root rot complex”. Among the pathogenic agents of the

root rot complex several species belonging to *Cylindrocarpon* sp., *Pythium* sp. *Rhizoctonia* sp. are known (Savory, 1966; Mazzola, 1998).

Call the replant problems affecting all fruit tree crops of high economic value in Europe are strongly related to land degradation problems due to intensive cropping with frequent orchard replants, this problem is particularly severe in Mediterranean countries where climate (cool humid winters alternating with warm dry summers) lead to a rapid decline in the soil organic matter (SOM) content such as Turkey. The main consequence of progressive SOM depletion is microbial biomass reduction and loss of microbial diversity, with reduction of soil functionality. Microbial diversity, along with microbial biomass, is the best indicator of soil health (Winding and Nielsen, 2002) and represents one of the main components of soil suppressiveness (Garbeva et al., 2004). This can be defined as the potential biological ability of soil to deal with soil-borne pathogens (Borneman and Becker, 2007). Bacterial communities have been widely used as a microbial indicator of soil diversity and soil functioning (Blackwood et al., 2007); however, recently soil fungi have been found to be a good indicator of functioning of agricultural soils (Suzuki et al., 2009).

The occurrence of replant disease is becoming an indicator of a general degraded state of fruit tree orchards. The only effective strategy to control replant disease and crop decline in organic fruit tree orchards is to increase the soil diversity, and consequently, the various microbial processes involved in controlling soil borne pathogens, enhancing root growth and mediating plant nutrition. (Suzuki et al., 2009).

The commercial market of bio-pesticides has expanded significantly in the last five to ten years. Some formulations of fungi and bacteria have been registered as biocontrol agents, but many bio-products have been marketed as plant growth promoters or plant strengtheners. They include inorganic compounds, composts, organic amendments, homeopathic compounds, microorganisms. These commercial products are not sufficiently regulated by legislation in several countries; in many cases they do not provide the name of biocontrol agents included in the formulation or do not quote the presence of microorganisms; moreover, there is a dearth of published scientific data. As far as the use of these compounds in Europe is concerned, many bio-pesticides come abroad and they contain microorganisms which actually represent alien organisms introduced into our environment. This last aspect has not yet been tackled by legislation, but risks deriving from the introduction of alien organisms should be considered in the medium term. Each of the above mentioned aspects, combined with the partial efficacy of commercial bio-pesticides and the variability of their effectiveness in controlling diseases (Whipps and Davies, 2001), have increased confusion and reduced the interest of producers in the biological control of soil borne pathogens. This could be performed acting on the soil autochthonous microflora, a naturally available resource, with a multi-factorial approach, by working with suitable cropping practices on the numerous biological components of soil suppressiveness.

The combination of the selected resources with low input agro-management practices for controlling soil borne pathogens, such as solarisation, bio-pesticides, soil movement and other available tools etc., should lead to new management options in organic fruit tree orchards. Many of these innovative management options could be easily transferred to other fruit crops such as stone fruits and strawberries, whose yield and quality decline due to the same biological components observed in fruit tree crops.

Phytophthora root rot and gummosis caused by *Phytophthora citrophthora* (Sm. and Sm.) Leonian are major problems in citrus orchards worldwide. This disease was first reported in Turkey in 1958, and it has affected lemon, mandarin, orange, grapefruit and sour orange cultivars (Salih, 1974)

Survey results in the 1970's the disease incidence in lemon orchards in the provinces of Içel and Adana to be 24% and 22.4%, respectively. Some cultural practices such as bud unions at soil surface level or less than 15 cm above soil line, overwatering, poor drainage, soil structure and high clay levels enhance the incidence of the disease. Methods of controlling *Phytophthora* root rot are generally preventative, e.g., cultural practices and fungicide application to trunk and lower branches. When infection does occur, removal of the infected bark is recommended; however, bark removal diminishes tree performance and fruit yield. This technique is also laborious and uneconomical when attempted on a large scale.

In present study was conducted to determine effect of some product of Bio-Formulates in citrus plants which protect from replant disease such as *Phytophthora citrophthora*.

2. MATERIALS AND METHODS

Soils were collected from fields which infected with *Phytophthora citrophthora*. Experiment, Sour orange and Trifliage rootstocks were used and treated ten active ingredient/microorganism for *Phytophthora citrophthora*. We used dose according to recommended company's dosage. Bio-formulates were used (Table 1). Sour orange and Trifliage rootstock plants were transplanted to a 15 liter pot containing to an infested soil mixture. Treatments were applied to plants with 3 replications and every replicate resulted from 6 plants in greenhouse.

Table 1. The List of Product of Bio-Formulates used for Citrus rootstocks.

Name and action	Active ingredient/microorganism	Company or origin
Companion	<i>Bacillus subtilis</i> GB03	Growth products
Cropset	<i>Lactobacillus acidophilus</i> , plant extract, <i>MnSO4</i> , <i>FeSO4</i> , <i>CuSO4</i>	Ant Agriculture (Improcrop EU)
Isr-2000	<i>Lactobacillus acidophilus</i> , plant extract, yeast extract, benzoic acid	Ant Agriculture (Improcrop EU)
Subtilex	<i>Bacillus subtilis</i> MBI 600	Bioglobal
T22 Planter Box	<i>T. harzianum</i> (Rifai Irk Krl-Agz 403)	Hasel Agriculture
Tricho plus	<i>T. harzianum</i>	Bioglobal
Alexin 95 PS	Fosfor pentaoksin (P2O5) 52%, Potasyum Oksit (K2O) 42%	Sumitoma
Combat Plus	Plant activators	Bioglobal
Endo Roots Souble (23,8%)	<i>Glomus intraradices</i> <i>Glomus aggregatum</i> <i>Glomus mosseage</i> <i>Glomus clarum</i> <i>Glomus monosporus</i> <i>Glomus deserticola</i> <i>Glomus brasilianum</i> <i>Glomus etunicatum</i> <i>Gigaspora margarita</i>	Bioglobal

Plants were incubated for 4 months in greenhouse, and were kept constantly wet. Pots were subjected them in dripping at 6 h. three times during week. At the end of the growing season plant height (cm), hlant height (cm), root dry weight (g) and shoot dry weight (g) were recorded for all trees in all experiments. At the conclusion of the experiment, results were subjected to analysis of variance according to software programme SAS-JUMP 7.0.

3. RESULTS AND DISCUSSIONS

3.1. Plant Height (cm)

At the end of the growing season, *Plant Height* were recorded for all trees in all experiments. The differences between average *Plant Height*, were found important statistically. And the highest differences were recorded from ISR200(31.4%). Endo Roots Soluble was the lowest one (3.6%). Also, T22 Planter Box took second high value (27.6%) (Table 2).

Table 2. Average Plant Height(cm)

		Plant Height (cm)		
	Name and action	Applied	Control	% Differences
1	Cropset	87.0	70.5	23.4 bc
2	Compain	81.5	69.7	17.0bc
3	Subtilex	70.7	66.0	7.1c
4	T22 Planter Box	91.5	71.7	27.7ac
5	Combat Plus	102.2	82.7	23.6a
6	ISR200	93.5	71.2	31.4ab
7	Alexin	78.5	78.5	0.0bc
8	Endo Roots Soluble	72.3	69.8	3.6bc
9	Endo Roots Soluble	82.8	78.3	5.7ac
	LSD(5%)	12.06		

There was significant differences in the same column between the averages ($P \leq 0.05$),

3.2 Trunk Diameter

Avarage trunk diameter measured at end of the growing season and showed Table 3. As obtained results from Product of Bio-Formulates, ISR200 was highest difference value (15.8%) with regard to trunk diameter. But Alexin got lowest value (0.6%).

Table 3. Average Trunk Diameter (mm)

		Trunk Diameter (mm)		
	Name and action	Applied	Control	% Differences
1	Cropset	10.0	9.8	1.9bc
2	Compain	10.1	9.7	4.2bc
3	Subtilex	9.9	9.8	1.4c
4	T22 Planter Box	11.6	10.2	14.0ac
5	Combat Plus	12.0	10.4	15.4ab
6	ISR200	11.6	10.0	15.8ac
7	Alexin	10.4	10.4	0.6ac
8	Endo Roots Soluble	11.2	9.6	16.5ac
9	Endo Roots Soluble	12.1	11.0	9.5a
	LSD(5%)	1.25		

There was significant differences in the same column between the averages ($P \leq 0.05$),

3.3 Root Dry Weight

Average Root Dry Weight calculated at end of the growing season after dried at 65 C in etuve and showed at Table 4. ISR200 had the highest difference when compared with control plants with 34.7% difference value. Compain had the lowest difference value(0.5%) with regard to root dry matter content.

Table 4. Root Dry Weight (g)

	Name and action	Root Dry Weight (g)		
		Applied	Control	% Differences
1	Cropset	44.2	41.7	6.0ab
2	Compain	33.8	33.7	0.5b
3	Subtilex	35.8	34.2	4.9ab
4	T22 Planter Box	43.0	33.3	29.0ab
5	Combat Plus	58.3	41.0	42.3a
6	ISR200	55.0	40.8	34.7a
7	Alexin	44.0	35.8	22.8ab
8	Endo Roots Soluble	52.5	40.0	31.3ab
9	Endo Roots Soluble	48.3	40.0	20.8ab
	LSD(5%)	12.85		

There was significant differences in the same column between the averages ($P \leq 0.05$),

3.4 Shoot Dry Weight

Shoot dry matter content calculated after harvest and dried under 65 C in etuve. As obtained result, ISR200 had the highest difference value (60%), but T22 Planter Box had the lowest value (4.7%) (Table 5).

Table 5. Shoot Dry Weight (g)

	Name and action	Shoot Dry Weight (g)		
		Applied	Control	% Differences
1	Cropset	51.7	40.0	29.2b
2	Compain	43.8	38.8	13.1b
3	Subtilex	43.3	38.3	13.0b
4	T22 Planter Box	45.0	43.0	4.7b
5	Combat Plus	64.2	57.0	12.6c
6	ISR200	60.0	37.5	60.0a
7	Alexin	51.0	40.8	24.9b
8	Endo Roots Soluble	50.8	37.5	35.6b
9	Endo Roots Soluble	33.3	20.8	60.0a
	LSD(5%)	10.86		

There was significant differences in the same column between the averages ($P \leq 0.05$)

4. CONCLUSIONS

Phytophthora root rot and gummosis caused by *Phytophthora citrophthora* (Sm. and Sm.) Leonian are major problems in citrus orchards worldwide. This problem formed as replant disease at citrus production at mediterranean region. There are lots of pesticide application to protect from *Phytophthora citrophthora*. But when pesticides were applied, soils and envioriment under thread.

Best solution for this problem is bio formulates. In present study, effect of nine different bio formulate which have sale by different companies were determined. All product had positive effect on plant growing, but ISR200 had highest value. This differences can be because of plant material, temperature and product doses.

5. ACKNOWLEDGEMENTS

Present study supported by EU CORE ORGANIC II Project

6. REFERENCES

- Savory, B.M. (1966). Specific replant diseases causing root necrosis and growth depression in perennial fruit and plantation crops. *Research Rev. Commonw. Bur. Horth. Pl. Crops* 64 pp
- Manici, L.M., Ciavatta, C., Kelderer, M., Erschbaumer, G. (2003) Replant problems in South Tyrol: role of fungal pathogens and microbial populations in conventional and organic apple orchards. *Plant & Soil* 256, 315-324
- Mazzola, M. (1998) Elucidation of the microbial complex having a causal role in the development of apple replant disease in Washington. *Phytopathology* 98, 930-938.
- van Schoor, L., S. Denman, Cook, N.C. (2009). Characterization of apple replant disease under South African conditions and potential biological management strategies. *Sci. Hort.* 119, 153
- Gordon Braun, P., Fuller K.D., McRae, K., Sherry, A.E. (2010) Fillmore Response of 'Honeycrisp ' Apple Trees to Combinations of Pre-plant Fumigation, Deep Ripping, and Hog Manure Compost Incorporation in a Soil with Replant Disease *HortScience* 45, 1702–1707
- Browne, G.T., Connell, J.H., Schneider, S.M. (2006) Almond replant disease and its management with alternative pre-plant soil fumigation treatments and rootstocks. *Plant Disease* 90, 869–876.
- Mazzola M. (1998) Elucidation of the microbial complex having a causal role in the development of apple replant disease in Washington. *Phytopathology* 98, 930-938.
- Winding, A., Nielsen, M. N.(2002) Microorganisms as Indicators of Soil Health. *Techn. Rep n.388*. 82 pp.
- Garbeva P., van Veen J.A., van Elsas J.D. (2004) Microbial diversity in soil: selection of microbial populations by plant and soil type and implications for disease suppressiveness. *Annual Review of Phytopathology*, 42, 243–70.
- Borneman, J., Becker, J.O. (2007) Identifying Microorganisms Involved in Specific Pathogen Suppression in Soil. *Ann. Rev. Phytopath.* 45, 153-172.
- Blackwood, C.B., Hudleston, D., Zak, D.R., Buyer, J. S. (2007). Interpreting Ecological Diversity Indices Applied to Terminal Restriction Fragment Length Polymorphism Data: Insights from Simulated Microbial Communities. *Appl. Environ. Microbiol.* 73, 5276-5283
- Suzuki, C., Nagaoka, K., Shimada, A., Takenaka, M. (2009) Bacterial communities are more dependent on soil type than fertilizer type, but the reverse is true for fungal communities. *Soil Sci. Pl Nutrition* 55, 80–90
- Manici, L. M., Caputo, F. (2010). Soil fungal communities as indicators for replanting new peach orchards in intensively cultivated areas. *European journal of agronomy*, 33(3), 188-196.
- Cook, R., Rayner, A.D.M. (1984) Ecology of Saprotrophic Fungi. Longman, London, New York, 415 pp. *European Fruit Summit* 2th ed. 2010 Cesena, Italy
- Whipps, J.M., Davies, K.G. (2001) Success in biological control of plant pathogens and nematodes by microorganisms, status and prospects. In Butt et al. Fungal as biological agents progress, problems and potential. 2001. 1-8. *CAB International* New York.